

Phytogenics support energy efficiency of ruminants

By Gonçalo Martins, Technical Services Lead Ruminants - EMEA

Cargill Micronutrition & Health Solutions

The efficiency of energy utilisation in ruminants codetermines the profitability of meat production, as well as its environmental impact. Through genetic selection and the development of new technologies and methodologies the efficiency of ruminants converting feed energy into meat for human consumption can be improved.

The efficiency of energy utilisation in ruminants is a major determinant of the profitability of meat production, as well as its environmental impact. This is why it has long been a concern for animal nutritionists and farmers. Through genetic selection and the development and application of new technologies and methodologies, animal scientists and producers have improved the efficiency of ruminants converting feed energy into meat for human consumption.

Energy efficiency



Ruminants are dependent upon microorganisms, through rumen fermentation, for digestion. During rumen fermentation, bacteria produce byproducts that can affect the efficiency of digestion. Furthermore, ruminants have a high maintenance requirement compared to other livestock (more than 50% of their feed intake).

When comparing meat production by cattle to other livestock, simple energetic efficiency is typically lower due to their reliance on forages and fibrous co-

products as feeds and the role of fermentation in their digestion.

Metabolic disorder: Acidosis



Feeding high-concentration diets favours beef growth, but an excessive amount of rapidly fermented starch could result in subacute ruminal acidosis (SARA). This intensive feeding system requires good control of ruminal fermentation, since beef cattle diets are very low on fiber, having an impact on chewing and ruminating activities and also saliva production, therefore reducing the physiological buffering capacity of the animal.

One of the biggest impacts of acidosis in beef cattle is the decrease in dry matter intake, which has a direct impact on the performance of the animals. This feed depression is linked with ruminal hypomotility by direct action of the high levels of volatile fatty acids (VFAs) since chemical receptors in the epithelium send a feedback signal to the brain to reduce ruminal motility.

In addition, severe liver abscesses, which result from acidosis, may reduce daily gain by 11% and feed efficiency by 9%. Also seen frequently associated with acidosis is diarrhea, usually along with undigested feed material. Impaired ruminal function in terms of rumination, bacterial breakdown, and a passage leads to changes in faecal aspects.

When formulating diets for beef cattle, we should consider the total amount of starch and the ruminal degradability of cereal grains. We can consider two types of starch sources according to the starch degradation rate :

- Fast degradable starch in grains and roots: cassava, wheat, barley, and triticale.
- Slowly degradable starch in grains and roots: rice, maize, sorghum, and potatoes.


Intestinal digestion of starch



As mentioned previously, starch from cereal grains is the primary dietary energy source for highly productive cattle. For most grains, 90% or more of the starch is normally fermented in the rumen. The starch that escapes fermentation and flows to the small intestine may be more resistant to enzymatic digestion. On average, 35% to 60% of starch entering the small intestine is degraded there.

However, when it comes to maize and sorghum, it's different. Due to a slower rate of starch digestion, up to 40% of maize starch can occasionally be found to escape fermentation in the rumen. This is important because the capacity of the ruminant small intestine to digest large amounts of starch has been questioned, as a consequence of the low levels of pancreatic amylase, intestinal maltase, and isomaltase but also because of low glucose absorption. Physical treatments impact the cost of diets but can increase starch fermentation in the rumen and digestibility in the intestine. The most frequent treatments are grinding, pelleting, dry or steamrolling, steam flaking, extrusion, popping, and fermentation.

Phytogenic feed solutions in beef cattle



Besides good management and a well-balanced diet, one of the ways to reduce feed costs and improve feed efficiency of feedlot cattle is to use phytogenic feed solutions. Substances such as saponins, essential oils, and spices have been shown to improve feed efficiency and, therefore, performance in beef cattle.

The unique formulation of Actifor Power can impact beef cattle performance since the phytogenic ingredients in its composition reduce the starch degradation rate in the rumen while stimulating saliva production. Combined, these two actions contribute to stabilising the physiological conditions of the rumen.

It can also help to optimise intestinal starch digestion, improve bypass starch utilisation, improve the energy efficiency of beef cattle diets, and reduce losses through feces.

	Control	Actifor™ Power
Feces consistency score⁽¹⁾	3.3	2.8
Fibers presence score⁽²⁾	3.8	3.6
Sieve residue (%)	37.5	29.0
Starch in feces (g/kg DM)	1.99	1.82

(1) 1 liquid à 5 hard (2) 1 short à 5 long

Table 1- Effect of Actifor™ Power on fiber and starch digestibility (Delacon in vivo trial, Germany 2015)

Table 1 illustrates the beneficial effects of this innovative phytogenic solution on fiber and starch digestibility parameters. The conversion of feed energy to the final product is an essential determinant of the profitability of meat production. One of the major sources of variation in the conversion of feed energy to net energy (NE) in beef cattle is the extent of feed digestibility. Thus, phytogenic feed solutions improve feed digestibility, and therefore feed conversion ratio (FCR, -0.07) and average daily gain (ADG, +114g) as shown in Figure 1.

Due to their positive impact on rumen function and animal welfare and performance, phytogenic feed solutions are expected to be a relevant tool in ruminant nutrition, featuring profitable natural animal production. When you choose a phytogenic solution from the Micronutrition and Health Solutions portfolio (part of the Cargill Animal Nutrition & Health business), you go beyond feed additives to incorporate technical expertise and digital tools to make holistic nutrition solutions.




Figure 1 – Actifor™ Power trial with 139 Charolais young bulls for 290 days (Italy, 2016)

This article is intended to provide scientific information only and should not be construed as marketing claims or guidance. The products discussed in this article are not available in all countries. Please contact a Delacon/Cargill representative to know more about product availability.

[Find out more about Micronutrition & Health Solutions](#)