



SOUTHER HEIFER MIN (RUM) (CN)

MEDICATED

A CONCENTRATE FOR DAIRY HEIFERS.

GROWING CATTLE ON PASTURE OR IN DRY LOT (STOCKER AND FEEDER, AND DAIRY AND BEEF REPLACEMENT HEIFERS) - For increased rate of weight gain.

Active Drug Ingredient

Monensin 716.53 g/ton

Guaranteed Analysis

Calcium (min.)	18.80	%	Potassium (min.)	0.80	%
Calcium (max.)	22.55	%	Selenium (min.)	9.00	ppm
Phosphorus (min.)	0.10	%	Manganese (min.)	1,164	ppm
Salt (min.)	11.65	%	Iodine (min.)	20	ppm
Salt (max.)	14.00	%	Cobalt (min.)	27	ppm
Sodium (min.)	4.75	%	Vitamin A (min.)	356,400	IU/lb.
Sodium (max.)	5.75	%	Vitamin E (min.)	590	IU/lb.
Magnesium (min.)	1.00	%			

Ingredients

Ground Limestone, Plant Protein Products, Grain Products, Salt, Magnesium Oxide, Potassium Sulfate, Magnesium Sulfate, Vitamin A Supplement, Soybean Oil, Vitamin D3 Supplement, Vitamin E Supplement, Zinc Amino Acid Complex, Sodium Selenite, Manganese Amino Acid Complex, Calcium Carbonate, Copper Amino Acid Complex, Manganous Oxide, Zinc Oxide, Basic Copper Chloride, Cobalt Glucoheptonate, Ferrous Sulfate, Potassium Iodide, Calcium Iodate, Mineral Oil, Cobalt Carbonate.

Tag Code: S3 181109 ~MTIME ~LOT

Feeding Directions

Feed at the rate of not less than 50 nor more than 200 mg per head per day in not less than 1 lb. of feed or, after the fifth day feed at the rate of 400 mg per head per day every other day in not less than 2 lb. of feed. During the first 5 days of feeding, cattle should receive no more than 100 mg per day contained in not less than 1 lb. of feed.

CAUTION: Do not allow horses or other equines access to feed containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not feed undiluted. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and the amount of refusals fed should be taken into consideration to prevent Monensin overdosing.

WARNING: A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

Custom Mix

For your safety, and that of your animals, always read and follow label directions and caution statements.



197904_38

MINERAL

722304459567



Net Weight Shown on Bag / Invoice
Cargill Animal Nutrition, P.O. Box 5614, Minneapolis, MN 55440